PUBLICATION REPORT

This document has been approved for public release and sale; its distribution is unlimited.

1560

42/89-90

ETHNIC DIFFERENCES IN THE PREVALENCE OF HEPATITIS DELTA AGENT IN DJIBOUTI

BY

E.A. Abbatte, Emile Fox, Said-Salah, Niel T. Constantine and Habiba H. Wassef

U.S. NAVAL MEDICAL RESEARCH UNIT NO.3 (CAIRO, ARAB REPUBLIC OF EGYPT) FPO NEW YORK 09527

56 **91-08717**

Short Report

Ethnic differences in the prevalence of hepatitis delta agent in Djibouti

E. A. Abbatte¹, Emile Fox^{2,3}, Said-Salah¹, Niel T. Constantine^{2,3} and Habiba H. Wassef⁴ Ministère de la Santé Publique, Djibouti; ²US Naval Medical Research Unit No. 3, Cairo, Egypt; ³International Health Program, University of Maryland School of Medicine, USA; ⁴World Health Organization, Djibouti.

Djibouti is a small country in the Horn of Africa, whose inhabitants belong to several ethnic groups. The north and south-west of the country is inhabited by Afars, an ethnic group different from the Somali Issas, who mostly live in the south-eastern part of the country. In addition, many people living in Djibouti originated from other countries (e.g. Ethiopia, Yemen, India, Europe).

Clinicians working in northern Djibouti have noticed for a long time that some young people might experience severe episodes of hepatic jaundice, which could lead to their premature death (Abbatte, unpublished observations). In southern Djibouti, however, such a phenomenon had never been recorded.

In October 1987 an extensive epidemiological serosurvey was conducted in Djibouti, involving mostly healthy young adults. Although the data collection was carried out in Djibouti city, situated in the south, the 656 subjects came from all over the country. Each subject had a medical interview and examination and his or her serum was tested for antibodies to various hepatitis viruses by commercially available enzymelinked immunosorbent assays (ELISA) (Abbott Laboratories, North Chicaco, USA). 11 sera were positive for antibodies to the delta agent (delta-Ab), all in healthy subjects also positive for hepatitis B virus (HBV) surface antigen. The prevalence of anti-delta antibodies was much higher in males than in females (3·2% versus 0·7% in the whole study population; P<0·02).

There was a striking difference in the prevalence of delta seropositivity according to ethnic groups. While the lowest infection rates were found in people originally from Ethiopia (2/231; 0.9%) and in people belonging to the Somali ethnic groups (2/204; 1%), the highest infection rates were recorded in the Afar people, since 6 of the 114 tested Afars were found delta-Ab positive (5.3%). All the positive Afar individuals were men, thus bringing the delta infection rate in male Afars to 7.7%, a value 10 times higher than the delta infection rate in the rest of the population combined (0.7%; P < 0.001). As shown in the Figure, the delta prevalence in HBsAg positive Afar males was 50%, much higher than the corres-

Correspondence should be addressed to Dr Emile Fox, US Naval Medical Research Unit No.3, F.P.O., New York, NY ponding rates for the combined non-Afars (10.5%; P<0.01).




Figure. HBsAg and delta antibody positivity according to the 3 major ethnic groups in Djibouti, the Afar people being divided according to sex. Column heights show the percentage prevalence rates in the general population of HBsAg positivity (whole column) and of delta agent in the HBsAg positive subjects (column base). The values under the 4 columns give the percentage delta prevalence in the HBsAg positive subjects.

In an attempt to study the mechanisms whereby the 6 Afar males might have been infected by the delta agent, we investigated them in more detail. However, no uniform transmission pattern could be found. The median age of the 6 subjects was 40 years; 2 individuals remembered a long-ago episode of severe jaundice; 5 had received medical injections; 3 had been in hospital, 2 of them for tuberculosis; 3 had previously had dental work, none surgery, and 1 had received a blood transfusion. 3 subjects admitted to extramarital sexual relations or to previous episodes of a sexually transmitted disease; none had scarifications or tattoos; none had current hepatomegaly or other liver-related abnormalities on clinical examination.

Delta agent is a unique hepatitis virus, since it requires hepatitis B virus (HBV) for its expression. It is transmitted among humans in ways similar to HBV. Although delta infection is thought to be world-wide, its geographical distribution appears irregular among countries where serosurveys have been performed. No satisfactory explanation exists to account for these discrepancies (RIZZETTO, 1983).

Delta agent can replicate in all primate species that support the replication of HBV, including man, chimpanzees and gorillas (RIZZETTO et al., 1983). Baboons live freely in Djibouti and it is interesting that they are more abundant in north Djibouti, where the Afar people live. Could it be that these baboons are a natural animal reservoir for the delta agent and a source of infection for the Afar people, in particular Afar males, who are more exposed to monkey bites since they are more mobile?

Future work on the delta agent in Djibouti will need to clarify its transmission and epidemiology and to assess the intimate relationship of delta infection with the acute and chronic liver diseases occurring in young Afar people.

^{09527-1600,} USA.

Requests for reprints should be addressed to Research Publications Division, US Naval Medical Research Unit No.3, F.P.O., New York, NY 09527-1600, USA.

Acknowledgements

We thank all medical and technical staff at the various institutions involved. This study was supported by the Naval Medical Research and Development Command, Naval Medical Command, National Capital Region, Bethesda, MD 20814, Work Units No. 3M161102BS10.AK.311 and No. 3M463105DH29.AA.002. The opinions and assertions contained herein are the private ones of the authors and are not to be construed as official or as reflecting the views of the Navy Department, the Naval Service at large, the Djibouti Ministry of Health or WHO.

References

Rizzetto, M. (1983). The delta agent. Hepatology, 3, 729-737.

Rizzetto, M., Verme, G. & Bonino, F. (editors) (1983). Viral Hepatitis and Delta Infection. New York: Alan R. Liss.

Received 24 May 1988; accepted for publication 23 August 1988

ROYAL SOCIETY OF TROPICAL MEDICINE AND HYGIENE

UNDERGRADUATE PROJECT PRIZE

The Royal Society of Tropical Medicine and Hygiene offers an annual prize of £200 for an account of work carried out by a non-medical undergraduate student of any nationality. The work will add to the knowledge of human or veterinary health or hygiene in the Tropics. Particular attention in the award of the prize will be directed towards originality and quality. It is intended that the prize should act as a stimulus for the pursuit of excellence in research carried out by undergraduates.

MEDICAL STUDENT ELECTIVE PRIZE

The Royal Society of Tropical Medicine and Hygiene offers an annual prize of £200 for an account of work carried out by a British medical student during an elective period spent in a tropical or developing country. In awarding this prize emphasis will be laid on the originality of the work and on

its contribution to knowledge or understanding of tropical diseases.

Further information and advice concerning both prizes can be obtained from the Honorary Secretaries, Royal Society of Tropical Medicine and Hygiene, Manson House, 26 Portland Place,

London, WIN 4EY.

	·		
Accesio	n For		
NTIS DTIC Unanno Justific	TAB Sunced	IN	QTIO
By Distribu	ution/	- 1	
A	vailability Code	. 1	
Dist	Avanila Char Special	L SAMESTON & A	
A-1	20	; ; ;	

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE						
:a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	16 RESTRICTIVE MARKINGS					
2a. SECURITY CLASS.F CATION AUTHORITY	3 DISTRIBUTION AVAILABILITY OF REPORT					
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE	Approved for public release: Distribution is unlimited.					
1 PERFORMING ORGANIZATION REPORT NUMBER(S)	5 MONITORING ORGANIZATION REPORT NUMBER(S)					
42/89-90						
U.S. Naval Medical Research Unit No. 3	7a NAME OF MONITORING ORGANIZATION					
Sc. ADDRESS (City, State, and ZIP Code)	7b. ADDRESS (City, State, and ZIP Code)					
PSC 452, Box 5000 FPO, AE 09835-0007						
3a. NAME OF FRONG: SPONSORING SD OFFICE-SYMBOL ORGANIZATION Naval Medical Re- (If applicable)	9. PROCUREMENT INSTRUMENT DENTIFICATION NUMBER					
search and Development Command NAVMEDRSCH	,					
3c. ADDRESS (City, State, and ZIP Code) National Naval Medical Center	10 SOURCE OF FU			IMORK LAUT		
Building 1, Tower 12	PROGRAM ELEMENT NO	PROJECT NO.3M4 31-	TASK NO	WORK UNIT ACCESSION NO		
Bethesda. MD 20889-5044	63105A	05H29	AA			
*1 TITLE (Include Security*Classification) Ethnic Differences in the Prevalence of Hejatitis Delta Agent in Djibouti. (UNCLASSIFIED). 12. PERSONAL AUTHOR(S) Abbatte. E.A.*, Fox, Emile, Said-Salah*, Constantine, Niel T. and Wassef. Habiba H.+						
	14. DATE OF REPORT (Year, Month, Day) 15 PAGE COUNT 1988. August, 23					
'6 SUPPLEMENTARY NOTATION						
Published in: Trans. R. Soc. Trop. Med. Hyg.,	83:107-108,	1989; Acc.	No. 1560.			
	Continue on reverse if necessary and identify by block number)					
FIELD GROUP SUB-GROUP Ethnic diff	erences; Hepatitis delta agent;					
Prevalence;	; Patients; Djibouti.					
As per attached. * Ministry of Public Health, Djibouti. * World Health Organization. Djibouti.						
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT 21 ABSTRACT SECURITY CLASSIFICATION						
☐ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS	UNCLASSIFIED 22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL					
Research Publications Branch	202-284-138		R.P.B.			